



Original research article

Sedentary behavior as a discriminator of functional disability among older adults: a cross-sectional study

Daniele Pereira de Santana¹, Lucas dos Santos^{1,2}, Italo Emmanoel Silva e Silva¹, Paulo Gonçalves Júnior¹, José Ailton Oliveira Carneiro¹, Thaís Alves Brito¹, Rafael Pereira³, Marcos Henrique Fernandes¹, Lucimere Bohn⁴, Jorge Mota⁴, Raildo da Silva Coqueiro^{1,4*}

¹ State University of Southwest Bahia, Department of Health, Center for Studies in Epidemiology of Aging, Jequié, Brazil

² State University of Tocantins, Health Science Complex, Augustinópolis, Brazil

³ State University of Southwest Bahia, Department of Biological Sciences, Integrative Physiology Research Center, Jequié, Brazil

⁴ University of Porto, Faculty of Sports, Research Centre of Physical Activity, Health, and Leisure, Laboratory for Integrative and Translational Research in Population Health, Porto, Portugal

Abstract

Objectives: To investigate the association between sedentary behavior and functional disability in instrumental activities of daily living among older adults, and to determine the accuracy and optimal cutoff point of sedentary behavior for the discrimination of functional disability.

Methods: This is a population-based, cross-sectional epidemiological survey of 318 older adults (mean age: 74.6 ± 9.7 years old) from Northeastern Brazil. Functional disability in instrumental activities of daily living was assessed using the Lawton scale (presence or absence). Sedentary behavior time was quantified using the International Physical Activity Questionnaire.

Results: The prevalence of functional disability was 59.7% (women: 68.5%; men: 48.5%; $P = 0.001$). The median sedentary behavior time was 4.7 h/day. Each 1 h/day increase in sedentary behavior was associated with a 4% increase in the likelihood of functional disability in women (PR: 1.04; 95% CI: 1.01–1.07). Furthermore, in women, sedentary behavior demonstrated an accuracy of 0.61 (95% CI: 0.53–0.68), a sensitivity of 23%, and a specificity of 100% (optimal cutoff point: 7.5 h/day).

Conclusion: Sedentary behavior was positively associated with functional disability in instrumental activities of daily living in women but not in men. Additionally, sedentary behavior showed accuracy in discriminating functional disability in women, with the optimal cutoff point identified at 7.5 h/day.

Keywords: Activities of daily living; Aging; Epidemiology; Sedentary lifestyle

Introduction

Aging is associated with an increased risk of morbidity (Santos et al., 2021, 2022a, b) and mortality (Santana et al., 2021). As a result of aging, detrimental structural changes occur in the neuromuscular system, such as reduction in the number of muscle fibers and, consequently in the cross-sectional area of skeletal muscle, as well as declines in excitatory impulses through supraspinal centers and in the recruitment capacity of motor units (Tieland et al., 2018). These changes significantly impact the ability of older adults to perform instrumental activities of daily living (IADL), leading to functional disability and compromising their quality of life (Tieland et al., 2018).

Functional disability among older adults represents a global public health challenge, albeit with substantial regional variation. In Europe, approximately 23.8% of adults aged

65 years and older have at least one IADL limitation (Portela et al., 2020), while in the Association of Southeast Asian Nations (ASEAN), pooled estimates indicate a considerably higher burden, with 46.8% of older adults experiencing IADL disability (Yau et al., 2022). In contrast, South Korea reports a lower prevalence of 15.5% for IADL limitations among adults aged 65 and older (Nguyen et al., 2022). In the United States, functional limitation is particularly pronounced among low-income working-age older adults, with IADL disability prevalence surpassing that of comparable European populations (Choi et al., 2024).

Concerning trends indicate that functional disability rates are increasing in several regions. In South Korea, a dramatic rise has been documented over a 12-year period, with IADL disability prevalence increasing from 10.7% in 2008 to 25.6% in 2020 (Nguyen and Hong, 2023). Similarly, Austria has experienced substantial increases between 2014 and 2019, with

* **Corresponding author:** Raildo da Silva Coqueiro, State University of Southwest Bahia, Department of Health, Av. José Moreira Sobrinho, s/n, Jequiezinho, Jequié – BA, Brazil, 45205-490; e-mail: rscoqueiro@uesb.edu.br
<http://doi.org/10.32725/kont.2026.009>

Submitted: 2025-10-12 • Accepted: 2026-01-28 • Prepublished online: 2026-01-29

KONTAKT 28/x: xxx-xxx • EISSN 1804-7122 • ISSN 1212-4117

© 2026 The Authors. Published by University of South Bohemia in České Budějovice, Faculty of Health and Social Sciences.

This is an open access article under the CC BY-NC-ND license.

IADL limitations rising from 18.9% to 35.1% in men, and from 38.2% to 50.8% in women (Woldemariam et al., 2024). These upward trends contrast with historical improvements observed in earlier birth cohorts and suggest a stalling or reversal of disability decline in cohorts born after 1945 in both the United States and Europe (Gimeno et al., 2024).

Future projections underscore the urgency of addressing functional disability. Microsimulation models for China predict that by 2048, nearly 38% of late middle-aged and older adults will be IADL-dependent (Jiang and Li, 2024). As populations continue to age globally, functional disability prevalence is expected to rise proportionally across most regions (United Nations, 2023), placing increasing demands on healthcare systems and caregiving resources worldwide.

A systematic review and meta-analysis estimated that the prevalence of IADL disability among Brazilian older adults was 43%, with regional analysis revealing a prevalence of 57% among older adults in the Northeast region (Meneguci et al., 2019). These figures substantially exceed the pooled estimate of 23.8% for Europe (Portela et al., 2020) and are comparable to the rate of 46.8% reported for the ASEAN region (Yau et al., 2022). This positions Brazil among the countries with the highest functional disability burden, particularly in socioeconomically disadvantaged regions. These high prevalence rates reflect an extremely adverse epidemiological panorama that parallels or exceeds international standards, as low functionality among older individuals has been associated with negative health outcomes such as depression, frailty syndrome (Aguiar et al., 2019), non-communicable chronic diseases, the need for emergency home care, and hospitalizations (Zanesco et al., 2020).

In addition to socioeconomic disadvantages, lifestyle and sociocultural patterns can influence daily activity and functional outcomes among older adults in Northeastern Brazil. National survey data show that the prevalence of limitations in activities of daily living is higher in the Northeast than in other Brazilian regions, particularly among women, reflecting broader gender and income inequalities in functional health status (Lima et al., 2020).

In this context, sedentary behavior that encompasses all activities performed at low energy expenditure (i.e., ≤ 1.5 metabolic equivalents), mostly performed in sitting and reclining postures (Tremblay et al., 2017), seems to prompt functional disability later in life (Scher et al., 2019). International evidence consistently demonstrates the association between sedentary behavior and functional disability. A prospective Japanese cohort study using accelerometer measurements found that greater sedentary time was associated with an increased risk of incident functional disability over nine years, with hazard ratios ranging from 1.21 to 1.45 for higher total sedentary time quartiles compared to the lowest quartile (Chen et al., 2023). In Brazil, national survey data indicate that television sedentary time ≥ 3 h/day is associated with 38% higher odds of IADL disability (Freitas et al., 2024). However, our literature search revealed only two studies identifying the accuracy of sedentary behavior as a discriminator of IADL disability among older adults (Brandão et al., 2019; Virtuoso Júnior et al., 2018). Notably, these two Brazilian studies employed Receiver Operating Characteristic (ROC) curve analysis to identify optimal sedentary behavior cutoff points. Their findings suggest that sedentary times of ≥ 603 min/day (Virtuoso Júnior et al., 2018) and ≥ 360 min/day (Brandão et al., 2019) increase the odds of IADL disability, demonstrating the potential utility of this methodological approach for establishing screening thresholds.

These findings highlight the need for more evidence on the discriminative power of sedentary behavior for IADL disability. Given the low cost and applicability of assessing sedentary behavior, the results of this research could inform health surveillance actions, enabling the early identification of older adults with functional impairments. Therefore, this study aimed to: (1) investigate the association between sedentary behavior and functional disability in IADL among community-dwelling older adults; (2) determine whether this association differs by sex; and (3) establish the discriminative accuracy and optimal cutoff point for daily sedentary behavior time in screening for IADL disability. Specifically, we sought to answer the following research questions: (1) Is sedentary behavior independently associated with IADL disability in older adults after controlling for relevant covariates? (2) Does the relationship between sedentary behavior and IADL disability differ between older men and women? (3) What is the optimal sedentary behavior threshold (h/day) that best discriminates older adults with and without IADL disability?

Material and methods

Study design, location, and population


This study was based on data from an epidemiological, cross-sectional, population-based, household survey entitled “Nutritional status, risk behaviors, and health conditions of older adults in Lafaiete Coutinho, Bahia, Brazil”. Details about the study site and population, as well as data collection methods, have been previously published (Queiroz et al., 2018). Briefly, a census was conducted in Lafaiete Coutinho to identify and include all older adults (≥ 60 years) residing in the urban area. Participants were recruited via the Family Health Strategy (FHS), which provides universal primary healthcare coverage across the municipality.

Individuals were included if they met the age (≥ 60 years) and residency (urban area) criteria. Exclusion criteria consisted of refusals to participate, individuals not located after three home visits on alternate days, and those with missing data for any of the variables of interest.

Of the 331 eligible older adults, 318 (96.1%) participated in the survey. There were 13 losses (3.9%), comprising three refusals (0.9%), and 10 individuals not found (3%) (Scheme 1).

Data collection

Data collection occurred in two stages. The first stage was conducted at participants’ homes and consisted of personal interviews, including sociodemographic information, health status, lifestyle factors, and motor performance tests. The second stage, which comprised anthropometry, was scheduled one to three days after the home interview and took place at the two



Scheme 1. Flowchart of the study participants

FHS units in the city. Before the interview, cognitive function was assessed using the modified Mini-Mental State Examination (MMSE). Older adults with scores <13 on the MMSE (Bertolucci et al., 1994) were asked to complete the Pfeffer scale; a score ≥ 6 indicated impaired cognitive function (Pfeffer et al., 1982). For participants with cognitive impairment, the help of a proxy respondent was requested for the interview.

Ethical considerations

This study was conducted in accordance with the Helsinki Declaration of the World Medical Association. It was approved by the Research Ethics Committee of the State University of Southwest Bahia (opinion: 491.661). The purpose and procedures of the study were explained to participants, who signed a Free and Informed Consent prior to data collection.

Functional disability in IADL (dependent variable)

Functional disability was assessed using Lawton's IADL scale (Lawton and Brody, 1969). Impairments were evaluated across six domains: preparing meals, managing money, going out, shopping, using the telephone, and taking medications. For each activity, participants were classified as independent (no help needed) or dependent (partial or total assistance needed). The IADL measure was then dichotomized as follows: 0 = independent (i.e., no limitations) and 1 = functional disability (i.e., difficulty performing at least one activity on the scale). This categorization is widely used in population-based studies to identify individuals at risk of losing autonomy (Aguilar et al., 2019; Brandão et al., 2019; Meneguci et al., 2019).

Sedentary behavior (independent variable)

The fifth domain of the International Physical Activity Questionnaire (IPAQ), which relates to sedentary behavior, was used (Craig et al., 2003). The IPAQ has been validated for Brazilian older adults (Benedetti et al., 2004, 2007). This questionnaire quantifies the time (h/day) spent sitting in various locations, including at home, in social settings, and at the doctor's office. This includes time spent sitting while resting, watching TV, visiting friends and relatives, reading, making phone calls, and eating, but excludes travel-related sitting time (e.g., on a bus or in a car). Based on this information, the weighted average for daily sedentary behavior was calculated using the following formula (Virtuoso Júnior et al., 2018): $\text{Time spent in sedentary behavior (h/day)} = [(\text{time spent sitting during weekdays} \times 5) + (\text{time spent sitting during weekend days} \times 2)] / 7$.

Confounders

The confounding variables included age group (60–69, 70–79, and ≥ 80 years); religion [yes = self-reported affiliation with any religion (e.g., Catholic, Protestant or Evangelical, Jewish, among others); no = no religious affiliation]; hospitalization in the last 12 months (yes and no); medication use (no use, 1 medication, and ≥ 2 medications); cognitive status (impaired and non-impaired); and nutritional status (low weight, adequate, and overweight). Cognitive status was assessed using the modified MMSE: <13 points = impaired; ≥ 13 points = not impaired (Bertolucci et al., 1994). Nutritional status was assessed as the body mass index [BMI: ratio of body mass to stature squared ($\text{BMI} = \text{kg}/\text{m}^2$)], and categorized as low weight ($\text{BMI} < 22 \text{ kg}/\text{m}^2$), adequate ($22 \text{ kg}/\text{m}^2 \leq \text{BMI} \leq 27 \text{ kg}/\text{m}^2$), and overweight ($\text{BMI} > 27 \text{ kg}/\text{m}^2$) (Gonçalves et al., 2019). Body mass and stature were measured as described by Pinheiro et al. (2020), using a portable digital scale (Zhongshan Camry Electronic, G-Tech Glass 6, China) and a portable compact sta-

diometer (Wiso, China), respectively. In 12 participants who could not remain standing, body mass and height could not be measured; therefore, these parameters were estimated using equations proposed by Chumlea et al. (1988).

Statistical analysis

Descriptive analysis of population characteristics included absolute and relative frequencies, means, medians, standard deviations (SD), and interquartile ranges (IQR). The normality of continuous data was verified using the Kolmogorov–Smirnov test. The chi-square and Mann–Whitney tests were employed to compare sex differences in the prevalence of functional disability in IADL and sedentary behavior time, respectively.

The association between functional disability in IADL and sedentary behavior was evaluated through Poisson regression with robust variance. To account for potential confounding factors, models stratified by sex were constructed to estimate prevalence ratios (PR) with their respective 95% confidence intervals (95% CI). Multivariable models were built using the backward method, where all confounding variables were initially included and subsequently removed one by one based on the highest *P*-values, up to a critical level of 0.10. Variables with a *P*-value ≤ 0.10 were retained in the final model for adjustment purposes, ensuring that the association between sedentary behavior and IADL disability was independent of sociodemographic and health-related factors.

The discriminative capacity of sedentary behavior for IADL disability was evaluated using parameters provided by the ROC curve (Nahm, 2022). First, the accuracy of sedentary behavior was examined through the area under the curve (AUC). Following this, the optimal cutoff point and its respective sensitivity and specificity values were determined using the Youden index (Martínez-Cambor and Pardo-Fernández, 2019). The significance level adopted in all analyses was 5% ($P \leq 0.05$). Data were analyzed using IBM SPSS Statistics for Windows (IBM SPSS, 21.0, 2012, Armonk, NY: IBM Corp.) and MedCalc (version 14.8.1, 2014).

Results

The study population consisted of 318 older adults (55.3% women), aged 60 to 108 years (mean and SD = 74.6 ± 9.7 years). Other characteristics of the population, by sex, are presented in Table 1. Men and women showed similarity regarding the distribution of age group, religion, and hospitalization. Women exhibited higher rates of using two or more medications, impaired cognitive status, and being overweight, while the proportions of non-use of medications and low weight were higher in men.



Overall, the prevalence of functional disability in IADL was 59.7%, and it was significantly higher in women compared to men (Chart 1).

The time (median and IQR) for sedentary behavior was 4.7 ± 4.4 hours, with no significant difference between sexes (Chart 2).

Functional disability in IADL was independently and positively associated with sedentary behavior in women but not in men. The results of the multiple regression model for women indicated that each one-unit (1 h/day) increment in sedentary behavior resulted in a 4% increase in the likelihood of functional disability in IADL, independently of age group, religion, hospitalization in the last 12 months, medication use, cognitive status, and nutritional status (Table 2).

Table 1. Characteristics of the study population by sex

Variables	% response	Sex	
		Female (n = 176)	Male (n = 142)
Age group, n (%)	100.0		
60–69 years		57 (32.3%)	48 (33.8%)
70–79 years		61 (34.7%)	60 (42.3%)
≥80 years		58 (33.0%)	34 (23.9%)
Religion, n (%)	99.7		
Yes		173 (98.3%)	134 (95.0%)
No		03 (1.7%)	07 (5.0%)
Hospitalization in the last 12 months, n (%)	99.7		
Yes		27 (15.3%)	29 (20.6%)
No		149 (84.7%)	112 (79.4%)
Use of medication, n (%)	96.9		
Does not use		25 (14.5%)	44 (32.6%)
1 medication		26 (15.0%)	24 (17.8%)
≥2 medications		122 (70.5%)	67 (49.6%)
Cognitive status, n (%)	93.7		
Impaired		64 (38.6%)	25 (18.9%)
Not impaired		102 (61.4%)	107 (81.1%)
Nutritional status, n (%)	95.3		
Low weight		37 (22.2%)	47 (34.6%)
Adequate		69 (41.3%)	62 (45.6%)
Overweight		61 (36.5%)	27 (19.9%)

**Chart 1.** Prevalence of functional disability in instrumental activities of daily living in older adults, according to sex. * Chi-square test**Chart 2.** Time of sedentary behavior of the older adults, according to sex. The box limits represent the first and third quartiles, while the internal line represents the median; the whiskers extend from the quartiles to the minimum and maximum values.

* Mann-Whitney test


Table 2. Prevalence ratios for functional disability in instrumental activities of daily living with increasing sedentary behavior in older adults, according to sex

Model	*PR	95% CI	P-value
Female	1.04	1.01–1.07	0.034
Male	1.00	0.93–1.07	0.994

Note: PR – prevalence ratio; CI – confidence interval. * Adjusted for age group (female) and for age group, medication use and cognitive status (male). The remaining confounders variables were removed from the models following the model criteria used in the study (see statistical analysis topic).

Because the regression analysis indicated that sedentary behavior was a predictor of functional disability in IADL only in women, the ROC curve analysis was performed to identify the cutoff point that best discriminates functional disability for this sex.

The AUC value indicates that sedentary behavior was able to distinguish older women with functional disabilities in IADL with an accuracy of 61% (Scheme 2). The cutoff point of 7.5 h/day of sedentary behavior was the best criterion to discriminate between older women with and without functional disabilities in IADL. Sensitivity was 23%, and specificity was 100%.

**Scheme 2.** ROC curve of sedentary behavior as a discriminator of functional disability in IADL in older women. AUC – area under the curve. IC – confidence interval.

Discussion

The results of the present study indicate that functional disability in IADL is independently and positively associated with sedentary behavior in older women, but not in older men. Additionally, the identified cutoff point of 7.5 h/day is the best criterion to discriminate functional disability in older women, presenting high specificity (100%) but low sensitivity (23%), which limits its utility for early screening purposes.

Some research conducted with populations from other countries has investigated the association between seden-

tary behavior and functional disability. For example, Chen et al. (2016) evaluated 1,634 older Japanese individuals using a triaxial accelerometer and showed that, regardless of sex, less sedentary time and a greater number of interruptions in sedentary time were related to a lower risk of IADL disability, even after adjusting for moderate to vigorous physical activity (MVPA). Dunlop et al. (2015) also used direct measurement (i.e., accelerometry) to examine the association between functional disability and sedentary time in a sample of 2,286 American older adults, concluding that, regardless of sex and time spent in MVPA, the presence of IADL disability was associated with daily sedentary hours and the percentage of daily sedentary waking hours.

These findings are partially in agreement with those of the present study, where sedentary behavior was positively associated with functional disability in older women but not in men. It is essential to highlight the methodological differences that may have influenced the results of the studies. The two aforementioned studies measured sedentary time directly (using motion sensors) (Chen et al., 2016; Dunlop et al., 2015), whereas we used an indirect measurement (questionnaire) that measures time spent in activities in the sitting position.

Furthermore, the inconsistent results across studies may have resulted from differences in analysis procedures. For example, in this research, the analyses were stratified by sex, whereas in the studies by Chen et al. (2016) and Dunlop et al. (2015), the analyses were adjusted for sex.

The sex-specific findings observed in our study – where sedentary behavior was significantly associated with IADL disability in women but not in men – warrant deeper interpretation considering both biological and sociocultural factors. From a biological perspective, older women experience more pronounced age-related declines in muscle mass and strength (sarcopenia) compared to men, particularly after menopause, due to declining estrogen levels. Estrogen exerts a protective effect on muscle tissues through estrogen receptors, and its deficiency accelerates muscle protein degradation and reduces muscle regenerative capacity (Santos et al., 2021; Tieland et al., 2018). Consequently, the deleterious effects of prolonged sedentary behavior on muscle metabolism and neuromuscular function may manifest more severely in women, making them more vulnerable to functional limitations. Additionally, women typically have lower baseline muscle mass and strength than men throughout the lifespan. This sexual dimorphism leaves them with less physiological reserve to withstand the adverse effects of sedentary behavior (Santana et al., 2021; Santos et al., 2022a). This lower threshold for functional impairment suggests that increased levels of sedentary behavior may trigger disability more rapidly in women than in men.

From a sociocultural perspective, gender differences in functional capacity and activity patterns among older adults have been consistently documented. Evidence from Brazilian population-based studies indicates that older women have a higher prevalence and earlier onset of functional limitations compared to men, which has been partly attributed to lifelong gendered roles and social inequalities (Barbosa et al., 2005). As functional limitations progress, reduced participation in domestic and social activities may contribute to greater sedentary behavior and social withdrawal, reinforcing sex-specific trajectories of disability in later life. Longitudinal evidence also suggests that women experience a faster transition to physical disability than men, highlighting the interaction between gender, functional decline, and activity patterns in aging populations (Mandal and Subaiya, 2025). These biological and sociocultural factors likely synergize to produce the

observed sex-specific associations, although further research with longitudinal designs and mixed-methods approaches is needed to fully elucidate these mechanisms.

The relationship between sedentary behavior and functional disability in older individuals is based on the adverse effects of prolonged sitting time on body composition, such as low muscle mass (Gianoudis et al., 2015) and high adiposity (Reid et al., 2018), which influence functionality. Moreover, hypokinesia can exacerbate declines in the capacity of muscle groups to maintain dynamic balance (Le Roux et al., 2022) and produce strength and power (Ramsey et al., 2021).

Therefore, excessive time spent in sedentary activities affects functional capacity. Sardinha et al. (2015) found that fewer than seven interruptions per hour in sedentary behavior resulted in twice the chances of disability in performing IADL among 371 older Portuguese individuals, regardless of the time spent in MVPA. Such findings reinforce the need for constant interruptions in hypokinetic behavior to improve performance among older adults (Santos et al., 2023a, b).

This study identified a cutoff point of 7.5 h/day of sedentary behavior that discriminates functional disability in IADL with an accuracy of 61% in women. While this cutoff demonstrated high specificity (100%), its low sensitivity (23%) limits its clinical utility for screening purposes. High specificity indicates that the cutoff is effective at correctly identifying older women without functional disability (i.e., low false-positive rate), which may be useful in resource-limited settings to avoid unnecessary interventions among those at low clinical risk. Therefore, while this cutoff may have some value in specific clinical contexts, such as confirming low risk in apparently healthy older women or as part of a multi-component assessment, it should not be relied upon as a primary screening instrument. For effective early detection and intervention, sedentary behavior assessment should be combined with other more sensitive indicators of functional decline, such as objective physical performance tests (e.g., gait speed, chair stand test) or comprehensive geriatric assessment tools that capture multiple domains of functional capacity.

Two other Brazilian studies conducted with older populations investigated the accuracy of sedentary behavior in discriminating functional disability in IADL. In the Southeast region of Brazil, Virtuoso-Júnior et al. (2018) verified in 624 older adults that times >600 min/day (i.e., 10 h/day) and >614 min/day (i.e., 10.2 h/day) were the best cutoff points for men and women, respectively. On the other hand, in the Northeast region of Brazil, Brandão et al. (2019) determined in a population-based survey of 310 older adults that the best values of sedentary behavior to identify functional disability were >330 min/day (i.e., 5.5 h/day) for men and >270 min/day (i.e., 4.5 h/day) for women.

Such differences may be due to the disparities in the populations studied. Virtuoso Júnior et al. (2018) investigated a medium-sized municipality with higher socioeconomic indicators than the population of the present study. While the socioeconomic profile of the population studied by Brandão et al. (2019) is similar to that of Lafaiete Coutinho, the former location exhibits a larger population and higher demographic density. This reinforces the importance of further research across different regions and urban centers of the country, so that the cutoff points related to sedentary behavior are tailored to the characteristics of each population. The cutoff point may differ in various populations.

An important consideration in interpreting our findings is the potential for reverse causality, which is inherent to the cross-sectional design of this study. While we observed an

association between higher sedentary behavior and increased likelihood of functional disability in older women, the temporal sequence and direction of this relationship cannot be established from our data. Longitudinal studies with repeated measurements of both sedentary behavior and functional status, ideally employing causal inference methods such as cross-lagged panel models or marginal structural models, are essential to disentangle these temporal relationships and establish whether reducing sedentary time can prevent or delay functional decline in older adults.

The use of indirect measurement to assess sedentary behavior can be considered another limitation of this study. Measurements performed using self-report instruments are subject to memory bias, especially among older adults. However, in cases where participants presented with cognitive impairment, the information was provided by a proxy respondent (caregiver or accompanying person), as described in the methodological procedures, to minimize recall bias and ensure data reliability. However, the use of proxy respondents introduces another potential source of measurement error, as proxies may have limited knowledge of the participants' habitual daily sedentary time and may interpret questions differently than the older adults themselves. Nevertheless, the IPAQ is a standard instrument that has been culturally adapted to measure the levels of physical activity and sedentary behavior of the Brazilian population (Benedetti et al., 2004, 2007). Moreover, the IPAQ demonstrates high reliability and moderate criterion validity compared to accelerometers (Craig et al., 2003).

Conversely, this study's strength lies in its census perspective, allowing for the evaluation of the population contingent of a small municipality in Northeast Brazil, which has low socioeconomic indicators and, therefore, lacks simple and low-cost measures for the identification of older adults in conditions of greater vulnerability within health surveillance. Furthermore, the study controlled for important confounders related to functional performance in IADL, such as cognitive status, nutritional status (BMI), polypharmacy, and religious affiliation. The inclusion of these variables strengthens the analysis, as they are factors that influence mobility and social engagement – particularly participation in religious rituals – which can contribute to reducing sitting time and preserving functional capacity.

Conclusion

The present investigation found that sedentary behavior is positively associated with functional disability to perform IADL in older women, but not in men. Furthermore, sedentary behavior shows accuracy in discriminating functional disability in older women, with a cutoff point of 7.5 h/day being the best criterion.

Author contributions

DPS, JAOC, TAB, MHF, RSC – Concept design; DPS, LS, IESS, JAOC, TAB, MHF, RSC – Collection, analysis, and interpretation of data; DPS, LS, IESS, PGJ, JAOC, TAB, RP, MHF, LB, JM, RSC – Preparing the draft of the study; DPS, LS, IESS, PGJ, JAOC, TAB, RP, MHF, LB, JM, RSC – Review and final approval.

Data availability

The datasets analyzed in the current study are not publicly available due to ethical restrictions related to participant privacy, but are available from the corresponding author upon reasonable request.

Acknowledgments

The authors would like to thank all the older adults who participated in this study, the Municipality of Lafaiete Coutinho, Bahia, Brazil, and all the researchers at the Center for Studies in Epidemiology of Aging.

Funding

This research received financial support from the Bahia State Research Foundation, through the Scientific Initiation Grant awarded to Santana DP (Process: 4017/2018).

Conflict of interest

The authors have no conflict of interest to declare.

References

1. Aguiar BM, Silva PO, Vieira MA, Costa FM, Carneiro JA (2019). Avaliação da incapacidade funcional e fatores associados em idosos. *Rev Bras Geriatr Gerontol* 22(2): e180163. DOI: 10.1590/1981-22562019022.180163.
2. Barbosa AR, Souza JM, Lebrão ML, Laurenti R, Marucci Mde F (2005). Functional limitations of Brazilian elderly by age and gender differences: data from SABE Survey. *Cad Saude Publica* 21(4): 1177–1785. DOI: 10.1590/s0102-311x2005000400020.
3. Benedetti TRB, Antunes PC, Rodriguez-Añez CR, Mazo GZ, Petroski EL (2007). Reprodutibilidade e validade do Questionário Internacional de Atividade Física (IPAQ) em homens idosos. *Ver Bras Med Esporte* 13(1): 11–16. DOI: 10.1590/S1517-86922007000100004.
4. Benedetti TRB, Mazo GZ, Barros MVG (2004). Aplicação do Questionário Internacional de Atividades Físicas para avaliação do nível de atividades físicas de mulheres idosas: validade concorrente e reprodutibilidade teste-reteste. *R Bras Ci Mov* 12(1): 25–34. DOI: 10.18511/rbcm.v12i1.538.
5. Bertolucci PHE, Brucki SMD, Campacci SR, Juliano Y (1994). O mini-exame do estado mental em uma população geral: impacto da escolaridade. *Arq Neuro-Psiquiatr* 52(1): 1–7. DOI: 10.1590/S0004-282X1994000100001.
6. Brandão TL, Lago KN, Rocha SV (2019). Sedentary behavior and functional disability in older adults with low economic status: Monidi Study. *Rev Bras Cineantropom Desempenho Hum* 21: e55944. DOI: 10.5007/1980-0037.2019v21e55944.
7. Chen T, Chen S, Honda T, Kishimoto H, Nofuji Y, Narazaki K (2023). Accelerometer-measured sedentary behavior and risk of functional disability in older Japanese adults: a 9-year prospective cohort study. *Int J Behav Nutr Phys Act* 20(1): 91. DOI: 10.1186/s12966-023-01490-6.
8. Chen T, Narazaki K, Haeuchi Y, Chen S, Honda T, Kumagai S (2016). Associations of sedentary time and breaks in sedentary time with disability in instrumental activities of daily living in community-dwelling older adults. *J Phys Act Health* 13(3): 303–309. DOI: 10.1123/jpah.2015-0090.
9. Choi H, Jivraj S, Fors S (2024). Functioning outcomes at older working ages by income groups: comparing the US with 11 European countries. *Innov Aging* 8(Suppl 1): 137. DOI: 10.1093/geroni/igae098.0442.
10. Chumlea WC, Guo S, Roche AF, Steinbaugh ML (1988). Prediction of body weight for the nonambulatory elderly from anthropometry. *J Am Diet Assoc* 88(5): 564–568.
11. Craig CL, Marshall AL, Sjöström M, Bauman AE, Booth ML, Ainsworth BE, et al. (2003). International physical activity questionnaire: 12-country reliability and validity. *Med Sci Sports Exerc* 35(8): 1381–1395. DOI: 10.1249/01.mss.0000078924.61453.fb.
12. Dunlop DD, Song J, Arntson EK, Semanik PA, Lee J, Chang RW, Hootman JM (2015). Sedentary time in US older adults associated with disability in activities of daily living independent of physical activity. *Phys Act Health* 12(1): 93–101. DOI: 10.1123/jpah.2013-0311.

13. Freitas BP, Cândido LM, Wagner KJP, Lacerda ACR, Mendonça VA, De Micheli R, et al. (2024). Sedentary behavior, obesity, and disabilities in community-dwelling older adults: analysis of the Brazilian National Health Survey 2019. *Healthcare (Basel)* 12(16): 1625. DOI: 10.3390/healthcare12161625.
14. Gianoudis J, Bailey CA, Daly RM (2015). Associations between sedentary behaviour and body composition, muscle function and sarcopenia in community-dwelling older adults. *Osteoporos Int* 26(2): 571–579. DOI: 10.1007/s00198-014-2895-y.
15. Gimeno L, Goisis A, Dowd JB, Ploubidis GB (2024). Cohort differences in physical health and disability in the United States and Europe. *J Gerontol B Psychol Sci Soc Sci* 79(8): gbae113. DOI: 10.1093/geronb/gbae113.
16. Gonçalves TJM, Horie LM, Gonçalves SEAB, Bacchi MK, Bailer MC, Barbosa-Silva TG, et al. (2019). Diretriz BRASPEN de terapia nutricional no envelhecimento. *BRASPEN J* 34(3): 1–68.
17. Jiang Y, Li L (2024). Projections of functional dependence among the late middle-aged and older population from 2018–2048 in China: a dynamic microsimulation. *Glob Health Res Policy* 9(1): 15. DOI: 10.1186/s41256-024-00357-y.
18. Lawton MP, Brody EM (1969). Assessment of older people: self-maintaining and instrumental activities of daily living. *Gerontologist* 9(3): 179–186.
19. Le Roux E, De Jong NP, Blanc S, Simon C, Bessesen DH, Bergouignan A (2022). Physiology of physical inactivity, sedentary behaviours and non-exercise activity: insights from the space bedrest model. *J Physiol* 600(5): 1037–1051. DOI: 10.1113/jp281064.
20. Lima ALB, Espelt A, Bosque-Prous M, Lima KC (2020). Gender differences in disability among older adults in the context of social gender and income inequalities: 2013 Brazilian Health Survey. *Rev Bras Epidemiol* 23: e200002. DOI: 10.1590/1980-549720200002.
21. Mandal K, Subaiya L (2025). Gender differences in disability among older persons in India: Evidence from the longitudinal ageing study in India (LASI). *Arch Gerontol Geriatr Plus* 2(3): 100183. DOI: 10.1016/j.aggp.2025.100183.
22. Martínez-Cambor P, Pardo-Fernández JC (2019). The Youden index in the generalized Receiver Operating Characteristic curve context. *Int J Biostat* 15(1): /j/ijb.2019.15.issue-1/ijb-2018-0060/ijb-2018-0060.xml. DOI: 10.1515/ijb-2018-0060.
23. Meneguci CAG, Meneguci J, Tribess S, Sasaki JE, Júnior JSV (2019). Incapacidade funcional em idosos brasileiros: Uma revisão sistemática e metanálise. *RBCEH* 16(3): 98–124. DOI: 10.5335/rbceh.v16i3.9856.
24. Nahm FS (2022). Receiver operating characteristic curve: Overview and practical use for clinicians. *Korean J Anesthesiol* 75: 25–36. DOI: 10.4097/kja.21209.
25. Nguyen VC, Hong GS (2023). Change in functional disability and its trends among older adults in Korea over 2008–2020: a 4-year follow-up cohort study. *BMC Geriatr* 23(1): 219. DOI: 10.1186/s12877-023-03867-5.
26. Nguyen VC, Moon S, Oh E, Hong GS (2022). Factors associated with functional limitations in daily living among older adults in Korea: a cross-sectional study. *Int J Public Health* 67: 1605155. DOI: 10.3389/ijph.2022.1605155.
27. Pfeffer RI, Kurosaki TT, Harrach CH, Jr., Chance JM, Filos S (1982). Measurement of functional activities in older adults in the community. *J Gerontol* 37(3): 323–329. DOI: 10.1093/geronj/37.3.323.
28. Pinheiro PA, Coqueiro RDS, Carneiro JAO, Correia TML, Pereira R, Fernandes MH (2020). Anthropometric indicators as screening tools for sarcopenia in older adult women. *Enferm Clin* 30(4): 269–274. DOI: 10.1016/j.enfcli.2018.12.010.
29. Portela D, Almada M, Midão L, Costa E (2020). Instrumental activities of daily living (iADL) limitations in Europe: an assessment of SHARE data. *Int J Environ Res Public Health* 17(20): 7387. DOI: 10.3390/ijerph17207387.
30. Queiroz RS, Faria LMA, Carneiro JAO, Coqueiro RS, Fernandes MH (2018). Age and mini-mental state examination score can predict poor-quality spirometry in the elderly: a cross-sectional study. *Clinics* 73: e374. DOI: 10.6061/clinics/2018/e374.
31. Ramsey KA, Rojer AGM, D'Andrea L, Otten RHJ, Heymans MW, Trappenburg MC, et al. (2021). The association of objectively measured physical activity and sedentary behavior with skeletal muscle strength and muscle power in older adults: A systematic review and meta-analysis. *Ageing Res Rev* 67: 101266. DOI: 10.1016/j.arr.2021.101266.
32. Reid N, Healy GN, Gianoudis J, Formica M, Gardiner PA, Eakin EE, et al. (2018). Association of sitting time and breaks in sitting with muscle mass, strength, function, and inflammation in community-dwelling older adults. *Osteoporos Int* 29(6): 1341–1350. DOI: 10.1007/s00198-018-4428-6.
33. Santana FM, Premaor MO, Tanigava NY, Pereira RMR (2021). Low muscle mass in older adults and mortality: A systematic review and meta-analysis. *Exp Gerontol* 152: 111461. DOI: 10.1016/j.exger.2021.111461.
34. Santos L, Almeida CB, Valença Neto PF, Silva RR, Santos IC, Casotti CA (2023a). Habitual physical activity and sedentary behavior as predictors of dynapenia in older adults: a cross-sectional study. *Sao Paulo Med J* 142(1): e2023070. DOI: 10.1590/1516-3180.2023.0070.R1.190523.
35. Santos L, Miranda CGM, Silva IES, Santos PHS, Brito TA, Fernandes MH, Carneiro JAO (2022a). Anthropometric indicators as predictors of dynapenia in postmenopausal women. *Motriz: Rev Educ Fis* 28: e10220001522. DOI: 10.1590/S1980-657420220001522.
36. Santos L, Miranda CGM, Souza TCB, Brito TA, Fernandes MH, Carneiro JAO (2021). Body composition of women with and without dynapenia defined by different cut-off points. *Rev Nutr* 34: e200084. DOI: 10.1590/1678-9865202134e200084.
37. Santos L, Silva RR, Santana PS, Valença Neto PF, Almeida CB, Casotti CA (2022b). Factors associated with dynapenia in older adults in the Northeast of Brazil. *J Phys Educ* 33(1): e3342. DOI: 10.4025/jphyseduc.v33i1.3342.
38. Santos L, Valença Neto PF, Pedreira RBS, Rocha RR, Galvão LL, Almeida CB, et al. (2023b) Association of physical activity combined with sedentary behavior with dynapenia in older adults. *J Phys Educ* 34: e-3430. DOI: 10.4025/jphyseduc.v34i1.3430.
39. Sardinha LB, Ekelund U, Santos L, Cyrino ES, Silva AM, Santos DA (2015). Breaking-up sedentary time is associated with impairment in activities of daily living. *Exp Gerontol* 72: 57–62. DOI: 10.1016/j.exger.2015.09.011.
40. Scher LML, Guarda FRB, Barros MVG, Chen Z, Anton SD (2019). Sedentary Time and Disability in Older Adults: A Systematic Review and Meta-Analysis. *J Aging Sci* 7: 206. DOI: 10.35248/2329-884.19.07.206.
41. Tieland M, Trouwborst I, Clark BC (2018). Skeletal muscle performance and ageing. *J Cachexia Sarcopenia Muscle* 9(1): 3–19. DOI: 10.1002/jcsm.12238.
42. Tremblay MS, Aubert S, Barnes JD, Saunders TJ, Carson V, Latimer-Cheung AE, et al. (2017). Sedentary Behavior Research Network (SBRN) – Terminology Consensus Project process and outcome. *Int J Behav Nutr Phys Act* 14(1): 75. DOI: 10.1186/s12966-017-0525-8.
43. United Nations (2023). Department of Economic and Social Affairs. *World Social Report 2023: Leaving no one behind in an ageing world*. New York: UN. [online] [cit. 2025-01-22]. Available from: <https://social.desa.un.org/publications/world-social-report-2023>
44. Virtuoso Júnior JS, Tribess S, Rocha SV, Sasaki JE, Aparecida Garcia C, Meneguci J, Romo-Pérez V (2018). Sedentary behavior as a predictor of functional disability in older adults. *Rev Bras Ativ Fis Saúde* 23: 1–7. DOI: 10.12820/rbafs.23e0010.
45. Woldemariam S, Stein VK, Haider S, Dorner TE (2024). Trends over time in the deficit of (instrumental) activities of daily living in the Austrian population aged 65 years and older: Results from the Austrian Health Interview Survey series. *Wien Klin Wochenschr* 136(17–18): 488–496. DOI: 10.1007/s00508-024-02388-4.

46. Yau PN, Foo CJE, Cheah NLJ, Tang KF, Lee SWH (2022). The prevalence of functional disability and its impact on older adults in the ASEAN region: a systematic review and meta-analysis. *Epidemiol Health* 44: e2022058. DOI: 10.4178/epih.e2022058.
47. Zanesco C, Bordin D, Santos CB, Fadel CB (2020). Dificuldade funcional em idosos brasileiros: um estudo com base na Pesquisa Nacional de Saúde (PNS-2013). *Ciênc Saúde Coletiva* 25(3): 1103–1118. DOI: 10.1590/1413-81232020253.19702018.