



Correction: Interaction of SAR inducers and nitrogen for key fodder quality parameters and fodder yield under leaf blight infection in winter sown oat

Ramandeep Kaur · Ashlesha Atri · Anuradha · Meenakshi Goyal

© The Author(s), under exclusive licence to Springer Nature B.V. 2026

Correction: Phytoparasitica (2025) 54:18
<https://doi.org/10.1007/s12600-025-01351-x>

The original version of this article unfortunately contained an incorrect Figure captions for both Figs. 5 and 6. Captions of the aforementioned figures should be inter-changed so it reads:

Figure 5. Correlation matrix heatmap between disease severity, SAR treatments and nutritional quality parameters of genotypes 1 (OL 14) and genotype 2 (OL 15). PDI- Percent disease index; GFY- green fodder yield; DM- dry matter; CP- crude protein; NDF- Neutral detergent fiber; ADF- acid detergent fiber. Pearson correlation coefficient within a column or row followed by (* or **) are significantly different at the level of significance $P \leq 0.05$ or $P \leq 0.01$, respectively.

Figure 6. Photographic presentation of treated and control plot.

Publisher's Note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

The original article can be found online at <https://doi.org/10.1007/s12600-025-01351-x>.

R. Kaur
Department of Plant Pathology, Punjab Agricultural University, Ludhiana, Punjab 141004, India

A. Atri (✉) · M. Goyal
Department of Plant Breeding and Genetics, Punjab Agricultural University, Ludhiana, Punjab 141004, India
e-mail: ashlesha-atr@pau.edu

Anuradha
Regional Research Station, Punjab Agricultural University, Kapurthala, Punjab 141004, India